

# A JUZVINSKIĬ ADDITION THEOREM FOR FINITELY GENERATED FREE GROUPS ACTIONS

LEWIS BOWEN & YONATAN GUTMAN

**ABSTRACT.** The classical *Juzvinskiĭ Addition Theorem* states that the entropy of an automorphism of a compact group decomposes along invariant subgroups. Thomas generalized the theorem to a skew-product setting. Using L. Bowen's *f*-invariant we prove the addition theorem for actions of finitely generated free groups on skew-products with compact totally disconnected groups or compact Lie groups (correcting an error from [Bo10c]) and discuss examples.

## CONTENTS

1. Introduction	1
Organization	2
Acknowledgements	2
2. The <i>f</i> -invariant	2
3. Skew-products	4
4. Relative entropy	6
5. A key Lemma	7
6. Proof of Theorem 3.3	8
7. Examples	11
7.1. The Ornstein-Weiss Example	11
7.2. A generalization	11
7.3. An algebraic example	12
References	14

## 1. INTRODUCTION

The following result was proven independently by H. Li [Li11] and Lind-Schmidt [LS09].

**Theorem 1.1.** *[Addition theorem for amenable groups] Let  $\Gamma$  be a countable discrete amenable group,  $G$  be a compact metrizable group and  $\alpha : \Gamma \rightarrow \text{Aut}(G)$  an action of  $\Gamma$  on  $G$  by group-automorphisms. Suppose  $N \triangleleft G$  is a closed normal  $\alpha(\Gamma)$ -invariant subgroup. Denote by  $\alpha_N : \Gamma \rightarrow \text{Aut}(N)$  and  $\alpha_{G/N} : \Gamma \rightarrow \text{Aut}(G/N)$  the induced actions and by  $\mu_G, \mu_N, \mu_{G/N}$*

---

*Date:* August 9, 2012.

*2010 Mathematics Subject Classification.* 37A35, 20E05.

*Key words and phrases.* Juzvinskiĭ Addition Theorem, *f*-entropy, Rokhlin-Abramov Addition Formula, finitely generated free groups.

the Haar probability measures on  $G, N$  and  $G/N$  respectively. Then the entropies of these actions satisfy:

$$h_{\mu_G}(\alpha) = h_{\mu_N}(\alpha_N) + h_{\mu_{G/N}}(\alpha_{G/N}).$$

In the case  $\Gamma = \mathbb{Z}$ , this result is due to Juzvinskiĭ [Ju65] from which it receives its name. The case  $\Gamma = \mathbb{Z}^d$  was proven in [LSW90]. Special cases were obtained by Miles [Mi08] and Björklund-Miles [BM09].

The paper [Bo10a] introduced a measure-conjugacy invariant, called the *f-invariant*, for probability-measure-preserving actions of finitely generated free groups. (Later a more general theory of sofic entropy was introduced in [Bo10b], of which we have little to say in the present article). In [Bo10c], a proof is claimed that the above addition formula extends to the case when  $\Gamma$  is a finitely generated free group, the entropy is replaced with the *f-invariant*, and  $G$  is either totally disconnected, a Lie group, or a connected abelian group (whenever the *f-invariant* is well-defined). However, there is an error in the proof. We prove here that the statement remains correct if either  $G$  is totally disconnected (and a mild additional hypothesis is satisfied) or  $G$  is a Lie group and the action is by smooth automorphisms. See the corrigendum [BG12] for other corrections to [Bo10c]. The main result is Theorem 2.2 below. We also prove a skew-product addition formula in Theorem 3.3 which may be of independent interest.

**Organization.** §2 reviews the *f-invariant* and states the main theorem; §3 reviews skew-products and proves Theorem 2.2 from Theorem 3.3. In §5 and §6 Theorem 3.3 is proven; §7 discusses examples, including the Ornstein-Weiss example.

**Acknowledgements.** We would like to thank Hanfeng Li for helpful conversations and the Fields Institute where some of the work for this project occurred. Y.G. would like to thank Benjamin Weiss, Eli Glasner and Jon Aaronson for helpful discussions. L.B. was partially supported by NSF grants DMS-0968762 and DMS-0954606.

## 2. THE *f*-INVARIANT

Let  $\Gamma = \langle s_1, \dots, s_r \rangle$  be a rank  $r$  free group. Let  $\alpha$  be a measure-preserving action of  $\Gamma$  on a standard probability space  $(X, \mathcal{B}_X, \nu)$ . We consider  $\alpha$  as a homomorphism from  $\Gamma$  to the group of automorphisms of  $(X, \mathcal{B}_X, \nu)$  and write  $\alpha_g$  for  $\alpha(g)$  ( $\forall g \in \Gamma$ ). Let  $\mathcal{P} = \{P_1, P_2, \dots\}$  be a countable partition of  $X$  into measurable subsets. The Shannon-entropy of  $\mathcal{P}$  is

$$H_\nu(\mathcal{P}) \triangleq - \sum_{P \in \mathcal{P}} \nu(P) \log(\nu(P)).$$

By convention  $0 \log(0) \triangleq 0$ . If  $\mathcal{P}, \mathcal{Q}$  are two partitions of  $X$  then their *join* is defined by  $\mathcal{P} \vee \mathcal{Q} \triangleq \{P \cap Q : P \in \mathcal{P}, Q \in \mathcal{Q}\}$ . If  $W \subset \Gamma$  is finite, we let  $\mathcal{P}^W \triangleq \bigvee_{w \in W} \alpha_w \mathcal{P}$ . Note that  $\alpha$  is only implicit in this notation. If  $H_\nu(\mathcal{P}) < \infty$  then define

$$\begin{aligned} F_\nu(\alpha, \mathcal{P}) &= (1 - 2r)H_\nu(\mathcal{P}) + \sum_{i=1}^r H_\nu(\mathcal{P} \vee \alpha_{s_i} \mathcal{P}) \\ f_\nu(\alpha, \mathcal{P}) &= \inf_{n \geq 0} F_\nu(\alpha, \mathcal{P}^{B(n)}) \end{aligned}$$

where  $B(n)$  is the ball of radius  $n$  centered at the identity with respect to the word metric. The partition  $\mathcal{P}$  is said to be *generating* (for the action  $\alpha$ ) if the smallest  $\alpha(\Gamma)$ -invariant  $\sigma$ -algebra containing  $\mathcal{P}$  is  $\mathcal{B}_X$  (up to sets of measure zero). In [Bo10a] it is shown that if  $\mathcal{P}, \mathcal{Q}$  are finite-entropy generating partitions then  $f_\nu(\alpha, \mathcal{P}) = f_\nu(\alpha, \mathcal{Q})$ . So we define the  $f$ -invariant of the action by  $f_\nu(\alpha) \triangleq f_\nu(\alpha, \mathcal{P})$  where  $\mathcal{P}$  is any finite-entropy generating partition for  $\alpha$ . If there does not exist a finite-entropy generating partition for  $\alpha$  then  $f_\nu(\alpha)$  is undefined.

It will be useful to have an alternative formulation of the  $f$ -invariant for which we need the following definitions. For  $g \in \Gamma$ , let  $h_\nu(\alpha_g, \mathcal{P})$  denote the entropy rate of  $\mathcal{P}$  with respect to the  $\mathbb{Z}$ -action generated by  $\alpha_g$ . To be precise,

$$h_\nu(\alpha_g, \mathcal{P}) = \lim_{n \rightarrow \infty} \frac{1}{2n+1} H \left( \bigvee_{i=-n}^n \alpha_g^i \mathcal{P} \right).$$

The *entropy* of the action  $\alpha_g$  is  $h_\nu(\alpha_g) = \sup_{\mathcal{P}} h_\nu(\alpha_g, \mathcal{P})$  where the supremum is over all finite measurable partitions  $\mathcal{P}$  of  $X$ . Define

$$\begin{aligned} F_\nu^*(\alpha, \mathcal{P}) &= (1-r)H_\nu(\mathcal{P}) + \sum_{i=1}^r h_\nu(\alpha_{s_i}, \mathcal{P}) \\ f_\nu^*(\alpha, \mathcal{P}) &= \inf_{n>0} F_\nu^*(\alpha, \mathcal{P}^{B(n)}). \end{aligned}$$

One of the main results of [Bo10c] is:

**Theorem 2.1.** *Let  $\alpha$  be a measure-preserving action of  $\Gamma$  on a standard probability space  $(X, \mathcal{B}_X, \nu)$ . Then for any finite-entropy generating partition  $\mathcal{P}$  for  $\alpha$ ,  $f_\nu(\alpha) = f_\nu^*(\alpha, \mathcal{P})$ .*

The main result of this paper is:

**Theorem 2.2.** *Let  $\Gamma = \langle s_1, \dots, s_r \rangle$  be a rank  $r$  free group,  $G$  be a compact metrizable group and  $\alpha : \Gamma \rightarrow \text{Aut}(G)$  an action of  $\Gamma$  on  $G$  by group-automorphisms. Suppose  $N \triangleleft G$  is a closed normal  $\alpha(\Gamma)$ -invariant subgroup. Denote by  $\alpha_N : \Gamma \rightarrow \text{Aut}(N)$  and  $\alpha_{G/N} : \Gamma \rightarrow \text{Aut}(G/N)$  the induced actions and by  $\mu_G, \mu_N, \mu_{G/N}$  the Haar probability measures on  $G, N$  and  $G/N$  respectively. Suppose there exists finite-entropy generating partitions for  $\alpha, \alpha_N, \alpha_{G/N}$  and one of the following hold.*

- (1)  *$N$  is totally disconnected and there exists a clopen finite-index normal subgroup  $N_0 \triangleleft N$  such that  $\{gN_0 : g \in N\}$  is a generating partition for  $\alpha_N$ .*
- (2)  *$G$  is a compact Lie group and the action  $\alpha$  is by smooth automorphisms.*

Then

$$f_{\mu_G}(\alpha) = f_{\mu_N}(\alpha_N) + f_{\mu_{G/N}}(\alpha_{G/N}).$$

**Remark 2.1.** The proof shows slightly more: if case (1) occurs and  $\alpha_{G/N}$  has a finite-entropy generating partition, then  $\alpha$  automatically has a finite-entropy generating partition. This follows from Lemmas 3.2 and 6.4. To be more precise, Lemma 3.2 shows that  $\alpha$  is measurably conjugate to a skew-product action of the form  $\alpha_{G/N} \times_\sigma \alpha_N$ . If  $\mathcal{P}$  is a finite-entropy generating partition for  $\alpha_{G/N}$  and  $\mathcal{Q} = \{gN_0 : g \in G\}$  is a generating partition for  $\alpha_N$  of the kind described in case (1) above, then Lemma 6.4 shows that  $\mathcal{P} \times \mathcal{Q}$  is generating for  $\alpha_{G/N} \times_\sigma \alpha_N$ . Because  $\mathcal{P}$  has finite-entropy and  $\mathcal{Q}$  is finite,  $\mathcal{P} \times \mathcal{Q}$  has finite entropy as required.

**Remark 2.2.** Suppose  $N$  as above is totally disconnected and  $N_0 \triangleleft N$  is a closed finite-index normal subgroup (the fact  $N_0$  is closed and finite-index implies  $N_0$  is clopen). Let  $M = \bigcap_{g \in \Gamma} \alpha_g N_0$ . Note  $M$  is  $\alpha(\Gamma)$ -invariant. Let  $\alpha_{G/M}, \alpha_{N/M}$  be the induced actions on  $G/M$  and  $N/M$  respectively. Let  $\mu_{G/M}, \mu_{N/M}$  be the respective Haar probability measures. Suppose that  $\alpha_{G/M}$  and  $\alpha_{G/N}$  admit finite-entropy generating partitions. Note that the clopen partition  $\{gN_0/M : g \in G\}$  is a finite generating partition for  $\alpha_{N/M}$ , as it is the image of the clopen generating partition  $\{gN_0 : g \in G\}$  under the continuous projection  $\pi : N \rightarrow N/M$ . So the theorem above implies

$$f_{\mu_{G/M}}(\alpha_{G/M}) = f_{\mu_{N/M}}(\alpha_{N/M}) + f_{\mu_{G/N}}(\alpha_{G/N}).$$

By the previous remark, this formula holds as long as  $\alpha_{G/N}$  admits a finite-entropy generating partition.

### 3. SKEW-PRODUCTS

The proof of Theorem 2.2 is based on a more general skew-product theorem of independent interest, the construction of which we recall next.

**Definition 3.1.** Let  $\Gamma$  be a group. Let  $(X, \mathcal{B}_X, \nu)$  be a Lebesgue space equipped with a  $\Gamma$ -action  $\alpha$  and  $G$  be a compact group with Borel  $\sigma$ -algebra  $\mathcal{B}$  and Haar measure  $\mu$ . Let  $\beta$  be a  $\Gamma$ -action by group-automorphisms on  $G$ . Let  $\sigma : \Gamma \times X \rightarrow G$  be a cocycle for  $\beta$  and  $\alpha$ , i.e.,  $\sigma$  is a measurable mapping so that for all  $g, h \in \Gamma, x \in X$

$$(3.1) \quad \sigma(gh, x) = (\beta_g \sigma(h, x)) \cdot \sigma(g, \alpha_h x).$$

Define the *skew-product action*  $\alpha \times_\sigma \beta$  of  $\Gamma$  on  $X \times G$  by:

$$(\alpha \times_\sigma \beta)_g(x, y) = (\alpha_g x, (\beta_g y) \cdot \sigma(g, x)) \quad (g \in \Gamma, x \in X, y \in G).$$

The connection between skew-product actions and the addition theorem is the following standard result (which we obtained from [Li11, Proof of Corollary 6.3]).

**Lemma 3.2.** Let  $\Gamma$  be a countable group,  $G$  be a compact metrizable group,  $\alpha : \Gamma \rightarrow \text{Aut}(G)$  an action of  $\Gamma$  on  $G$  by group-automorphisms and  $N \triangleleft G$  a closed normal  $\alpha(\Gamma)$ -invariant subgroup. Denote by  $\alpha_N : \Gamma \rightarrow \text{Aut}(N)$  and  $\alpha_{G/N} : \Gamma \rightarrow \text{Aut}(G/N)$  the induced actions. Then there is a cocycle  $\sigma : \Gamma \times G/N \rightarrow N$  such that  $\alpha_{G/N} \times_\sigma \alpha_N$  is measurably conjugate with  $\alpha$ .

The main technical result of this paper is:

**Theorem 3.3.** Let  $\Gamma = \langle s_1, \dots, s_r \rangle$  be a rank  $r$  free group,  $\alpha$  a measure-preserving action of  $\Gamma$  on a standard probability space  $(X, \mathcal{B}_X, \nu)$ ,  $G$  a compact metrizable group,  $\beta$  an action of  $\Gamma$  on  $G$  by group-automorphisms, and  $\sigma : \Gamma \times X \rightarrow G$  a cocycle for these actions. Suppose that  $G$  is totally disconnected and there exists a finite-index clopen normal subgroup  $N \triangleleft G$  such that  $\{gN : g \in G\}$  is a generating partition for  $\beta$ . Let  $\mu$  denote the Haar probability measure on  $G$ . Suppose also that there is a finite-entropy generating partition for  $\alpha$ . Then

$$f_{\nu \times \mu}(\alpha \times_\sigma \beta) = f_\nu(\alpha) + f_\mu(\beta).$$

The analog of this theorem for discrete countable amenable groups  $\Gamma$  when  $G$  is an arbitrary compact metrizable group was established in [Li11]. The case  $\Gamma = \mathbb{Z}$  was proven earlier by Thomas [Th71] and the case  $\Gamma = \mathbb{Z}^d$  is shown in [LSW90].

Theorem 3.3 is proven in the next section. Next we combine this result with the following two lemmas to complete the proof of Theorem 2.2.

**Lemma 3.4.** *Let  $M$  be a smooth compact Riemannian manifold. Let  $T : M \rightarrow M$  be a diffeomorphism. Then  $h_\mu(T) < \infty$  for any  $T$ -invariant probability measure  $\mu$ .*

*Proof.* This is due to Kushnirenko [Ku65]. Alternatively, it follows from Ruelle's inequality (see e.g. [KH95, Corollary S.2.17]).  $\square$

**Lemma 3.5.** *Let  $\Gamma = \langle s_1, \dots, s_r \rangle$  be a rank  $r$  free group with  $r > 1$ ,  $M$  be a smooth compact Riemannian manifold,  $\alpha$  a measure-preserving action of  $\Gamma$  on  $M$  by diffeomorphisms and  $\mu$  a non-atomic  $\alpha(\Gamma)$ -invariant probability measure on  $M$ . Then  $f_\mu(\alpha) = -\infty$  if there is a finite-entropy generating partition for the action.*

*Proof.* Let  $m = \max_{i=1}^r h_\mu(\alpha_{s_i})$ . By the previous lemma,  $m < \infty$ . Let  $\mathcal{P}$  be a finite-entropy generating partition for  $\alpha$ . Let  $N > 0$ . Because  $\mu$  is non-atomic, there is a finite partition  $\mathcal{Q}$  of  $M$  with  $H_\mu(\mathcal{Q}) > N$ . So after replacing  $\mathcal{P}$  with  $\mathcal{P} \vee \mathcal{Q}$  if necessary, we may assume that  $H_\mu(\mathcal{P}) > N$ . By Theorem 2.1

$$\begin{aligned} f_\mu(\alpha) &= f_\mu^*(\alpha, \mathcal{P}) = \inf_{n>0} F_\mu^*(\alpha, \mathcal{P}^{B(n)}) \\ &\leq (1-r)H_\mu(\mathcal{P}) + \sum_{i=1}^r h_\mu(\alpha_{s_i}, \mathcal{P}) \\ &\leq (1-r)N + rm. \end{aligned}$$

Because  $N > 0$  is arbitrary and  $r > 1$ , this implies the lemma.  $\square$

*Proof of Theorem 2.2 from Theorem 3.3.* Suppose item (1) holds. By Lemma 3.2,  $\alpha$  is measurable conjugate with  $\alpha_{G/N} \times_\sigma \alpha_N$  for some cocycle  $\sigma$ . So Theorem 3.3 implies

$$f_{\mu_G}(\alpha) = f_{\mu_G}(\alpha_{G/N} \times_\sigma \alpha_N) = f_{\mu_{G/N}}(\alpha_{G/N}) + f_{\mu_N}(\alpha_N)$$

as required.

Suppose that item (2) holds; i.e.,  $G$  is a compact Lie group and  $\alpha$  is an action by smooth group-automorphisms. If  $G$  is finite then the theorem is clear because

$$\begin{aligned} f_{\mu_G}(\alpha) &= -(r-1) \log |G| = -(r-1) \log |G/N| - (r-1) \log |N| \\ &= f_{\mu_{G/N}}(\alpha_{G/N}) + f_{\mu_N}(\alpha_N). \end{aligned}$$

By Theorem 1.1, we may assume, without loss of generality, that  $r > 1$ . If  $G$  is infinite then, because it is compact, it has positive dimension. So  $\mu_G$  is non-atomic and the previous lemma implies  $f_{\mu_G}(\alpha) = -\infty$ . Also, because  $G$  is infinite, either  $N$  or  $G/N$  is infinite. Therefore, either  $\mu_N$  or  $\mu_{G/N}$  is non-atomic. Of course, the actions  $\alpha_N$  and  $\alpha_{G/N}$  are smooth (because every continuous homomorphism between Lie groups is analytic [He01, Ch. II, Theorem 2.6]). It should be noted that the  $f$ -invariant does not take on the value  $+\infty$ . So the previous lemma implies  $f_{\mu_{G/N}}(\alpha_{G/N}) + f_{\mu_N}(\alpha_N) = -\infty$ .  $\square$

#### 4. RELATIVE ENTROPY

The proof of Theorem 3.3 uses the relative  $f$ -invariant theory developed in [Bo10c], which we review here. Let  $(X, \mathcal{B}_X, \nu)$  be a standard probability space. Let  $\mathcal{P}$  be a countable measurable partition of  $X$  and let  $\mathcal{F} \subset \mathcal{B}_X$  be a sub-sigma algebra. Recall that for a.e.  $x \in X$ , the conditional expectation  $\mathbb{E}[\cdot|\mathcal{F}](x)$  is a probability measure on  $(X, \mathcal{B}_X)$  satisfying

- (1)  $x \mapsto \mathbb{E}[A|\mathcal{F}](x)$  is  $\mathcal{F}$ -measurable for any  $A \in \mathcal{B}_X$ ;
- (2)  $\int \mathbb{E}[A|\mathcal{F}](x) d\nu(x) = \nu(A)$  for any  $A \in \mathcal{B}_X$ .

The information function  $I(\mathcal{P}|\mathcal{F})$  is a function on  $X$  defined by

$$I(\mathcal{P}|\mathcal{F})(x) = -\mathbb{E}[P_x|\mathcal{F}](x) \log(\mathbb{E}[P_x|\mathcal{F}](x))$$

where  $P_x \in \mathcal{P}$  is the unique partition element with  $x \in P_x$ . The Shannon entropy of  $\mathcal{P}$  *relative to*  $\mathcal{F}$  is

$$H_\nu(\mathcal{P}|\mathcal{F}) = \int I(\mathcal{P}|\mathcal{F})(x) d\nu(x).$$

If  $T$  is a measure-preserving transformation of  $(X, \mathcal{B}_X, \nu)$  then the entropy rate of  $(T, \mathcal{P})$  *relative to*  $\mathcal{F}$  is

$$h_\nu(T, \mathcal{P}|\mathcal{F}) = \lim_{n \rightarrow \infty} \frac{1}{2n+1} H_\nu \left( \bigvee_{i=-n}^n T^i \mathcal{P} | \mathcal{F} \right).$$

This is well-defined whenever  $\mathcal{F}$  is  $T$ -invariant. We also define the entropy rate of  $T$  *relative to*  $\mathcal{F}$  by

$$h_\nu(T|\mathcal{F}) = \sup_{\mathcal{P}} h_\nu(T, \mathcal{P}|\mathcal{F})$$

where the supremum is over all finite-entropy partitions  $\mathcal{P}$  of  $X$ .

Now suppose  $\Gamma = \langle s_1, \dots, s_r \rangle$  and  $\alpha$  is a measure-preserving action of  $\Gamma$  on  $(X, \mathcal{B}_X, \nu)$ . Define

$$\begin{aligned} F_\nu(\alpha, \mathcal{P}|\mathcal{F}) &= (1 - 2r)H_\nu(\mathcal{P}|\mathcal{F}) + \sum_{i=1}^r H_\nu(\mathcal{P} \vee \alpha_{s_i} \mathcal{P}|\mathcal{F}) \\ f_\nu(\alpha, \mathcal{P}|\mathcal{F}) &= \inf_{n \geq 0} F_\nu(\alpha, \mathcal{P}^{B(n)}|\mathcal{F}). \end{aligned}$$

Also define

$$\begin{aligned} F_\nu^*(\alpha, \mathcal{P}|\mathcal{F}) &= (1 - r)H_\nu(\mathcal{P}|\mathcal{F}) + \sum_{i=1}^r h_\nu(\alpha_{s_i}, \mathcal{P}|\mathcal{F}) \\ f_\nu^*(\alpha, \mathcal{P}|\mathcal{F}) &= \inf_{n \geq 0} F_\nu^*(\alpha, \mathcal{P}^{B(n)}|\mathcal{F}). \end{aligned}$$

**Theorem 4.1.** *Let  $\alpha$  be a measure-preserving action of  $\Gamma$  on a standard probability space  $(X, \mathcal{B}_X, \nu)$ . If  $\mathcal{P}, \mathcal{Q}$  are any two finite-entropy generating partitions for  $\alpha$  and  $\mathcal{F} \subset \mathcal{B}_X$  is an  $\alpha(\Gamma)$ -invariant sub- $\sigma$ -algebra then  $f_\nu(\alpha, \mathcal{P}|\mathcal{F}) = f_\nu(\alpha, \mathcal{Q}|\mathcal{F}) = f_\nu^*(\alpha, \mathcal{P}|\mathcal{F}) = f_\nu^*(\alpha, \mathcal{Q}|\mathcal{F})$ .*

*Proof.* This is implied by [Bo10c, Theorems 5.3, 9.1]. The proof requires a small correction; see [BG12].  $\square$

Because of this theorem, we define  $f_\nu(\alpha|\mathcal{F}) \triangleq f_\nu(\alpha, \mathcal{P}|\mathcal{F})$  where  $\mathcal{P}$  is any finite-entropy generating partition for  $\alpha$ . If there does not exist a finite-entropy generating partition for  $\alpha$  then  $f_\nu(\alpha|\mathcal{F})$  is undefined.

**Theorem 4.2.** *[The  $f$ -invariant Abramov-Rokhlin Addition Formula] Let  $\alpha$  be a measure-preserving action of  $\Gamma$  on a standard probability space  $(X, \mathcal{B}_X, \nu)$ . Let  $\mathcal{P}, \mathcal{Q}$  be finite-entropy partitions of  $X$ . Let  $\Sigma(\mathcal{Q})$  be the smallest  $\Gamma$ -invariant sub- $\sigma$ -algebra containing  $\mathcal{Q}$ . Then*

$$f_\nu(\alpha, \mathcal{P} \vee \mathcal{Q}) = f_\nu(\alpha, \mathcal{Q}) + f_\nu(\alpha, \mathcal{P}|\Sigma(\mathcal{Q})).$$

*Proof.* This is [Bo10c, Theorem 1.3]. The proof requires a small correction; see [BG12].  $\square$

## 5. A KEY LEMMA

The purpose of this section is to prove the key lemma below for skew-products of  $\mathbb{Z}$ -actions. Let  $(X, \mathcal{B}_X, \nu)$  be a Lebesgue space,  $T \in \text{Aut}(X, \mathcal{B}_X, \nu)$ ,  $G$  a compact metrizable group, equipped with Haar measure  $\mu$  and  $S$  a group-automorphism of  $G$ . A cocycle for  $T$  and  $S$  is a cocycle for the actions of  $\mathbb{Z}$  induced by  $T$  and  $S$ . That is, it is a measurable map  $\sigma : \mathbb{Z} \times X \rightarrow G$  such that

$$(5.1) \quad \sigma(n+m, x) = (S^n \sigma(m, x)) \cdot \sigma(n, T^m x).$$

**Lemma 5.1.** *Let  $(X, \mathcal{B}_X, \nu), G, T, S, \sigma$  be as above. Let  $\mathcal{Q}$  be a finite-entropy partition of  $G$ . Let*

$$K(\mathcal{Q}) = \sup_{g \in G} H(\mathcal{Q}g|\mathcal{Q}) + H(\mathcal{Q}|\mathcal{Q}g).$$

*Then*

$$\left| h_{\nu \times \mu}(T \times_\sigma S, X \times \mathcal{Q}|\mathcal{B}_X) - h_\mu(S, \mathcal{Q}) \right| \leq K(\mathcal{Q}).$$

*Proof.* By the definition of conditional entropy :

$$\begin{aligned} & h_{\nu \times \mu}(T \times_\sigma S, X \times \mathcal{Q}|\mathcal{B}_X) \\ &= \lim_{m \rightarrow \infty} \frac{1}{m} H_{\nu \times \mu} \left( \bigvee_{k=0}^{m-1} (T \times_\sigma S)^{-k} X \times \mathcal{Q} | \mathcal{B}_X \right) \\ &= \lim_{m \rightarrow \infty} \frac{1}{m} \int H_{\delta_x \times \mu} \left( \bigvee_{k=0}^{m-1} (T \times_\sigma S)^{-k} X \times \mathcal{Q} \right) d\nu(x) \end{aligned}$$

where  $\delta_x$  is the Dirac probability measure concentrated on  $\{x\}$ .

We claim that for any set  $P \subset G$ ,

$$\{x\} \times G \cap (T \times_\sigma S)^{-k}(X \times P) = \{x\} \times S^{-k}(P\sigma(k, x)^{-1}).$$

Indeed,  $(x, y)$  is contained in  $(T \times_\sigma S)^{-k}(X \times P)$  if and only if

$$(T \times_\sigma S)^k(x, y) = (T^k x, (S^k y)\sigma(k, x)) \in X \times P$$

which occurs if and only if

$$y \in S^{-k}(P\sigma(k, x)^{-1}).$$

So if

$$\mathcal{Q}_x^m = \bigvee_{k=0}^{m-1} S^{-k}(\mathcal{Q}\sigma(k, x)^{-1}).$$

then

$$H_{\delta_x \times \mu} \left( \bigvee_{k=0}^{m-1} (T \times_\sigma S)^{-k} X \times \mathcal{Q} \right) = H_\mu(\mathcal{Q}_x^m)$$

which implies:

$$(5.2) \quad h_{\nu \times \mu}((T \times_\sigma S), X \times \mathcal{Q} | \mathcal{B}_X) = \lim_{m \rightarrow \infty} \frac{1}{m} \int_X H_\mu(\mathcal{Q}_x^m) d\nu(x)$$

Define:

$$\mathcal{Q}^m = \bigvee_{k=0}^{m-1} S^{-k} \mathcal{Q}$$

By the definition of entropy:

$$(5.3) \quad h_\mu(S, \mathcal{Q}) = \lim_{m \rightarrow \infty} \frac{1}{m} \int_X H_\mu(\mathcal{Q}^m) d\nu(x)$$

Note  $|H_\mu(\mathcal{Q}^m) - H_\mu(\mathcal{Q}_x^m)| \leq H_\mu(\mathcal{Q}^m | \mathcal{Q}_x^m) + H_\mu(\mathcal{Q}_x^m | \mathcal{Q}^m)$ . Thus:

$$\begin{aligned} |H_\mu(\mathcal{Q}^m) - H_\mu(\mathcal{Q}_x^m)| &\leq \sum_{k=0}^{m-1} H_\mu(S^{-k} \mathcal{Q} | S^{-k}(\mathcal{Q}\sigma(k, x)^{-1})) + H_\mu(S^{-k}(\mathcal{Q}\sigma(k, x)^{-1}) | S^{-k} \mathcal{Q}) \\ &= \sum_{k=0}^{m-1} H_\mu(\mathcal{Q} | \mathcal{Q}\sigma(k, x)^{-1}) + H_\mu(\mathcal{Q}\sigma(k, x)^{-1} | \mathcal{Q}) \leq mK(\mathcal{Q}). \end{aligned}$$

Finally (5.2) and (5.3) imply  $|h_{\nu \times \mu}((T \times_\sigma S), X \times \mathcal{Q} | \mathcal{B}_X) - h_\mu(S, \mathcal{Q})| \leq K(\mathcal{Q})$ .  $\square$

## 6. PROOF OF THEOREM 3.3

For this section, let  $\Gamma, (X, \mathcal{B}_X, \nu), (G, \mathcal{B}_G, \mu), \alpha, \beta, \sigma$  be as in Theorem 3.3. A *special partition* of  $G$  is a partition  $\mathcal{Q}$  such that there exists a finite-index normal clopen subgroup  $N < G$  such that  $\mathcal{Q} = \{gN : g \in G\}$ . The next lemma is left as an exercise for the reader.

**Lemma 6.1.** *If  $\mathcal{Q}$  is special and  $T_1, \dots, T_n$  are automorphisms of  $G$  then  $\bigvee_{i=1}^n T_i \mathcal{Q}$  is also special.*

**Lemma 6.2.** *If  $\mathcal{P}$  is any finite-entropy partition of  $X$  and  $\mathcal{Q}$  is a special partition of  $G$  then*

$$F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, \mathcal{P} \times \mathcal{Q} | \mathcal{B}_X) = F_\mu^*(\beta, \mathcal{Q}).$$



*Proof.* Because  $\mathcal{Q}g = \mathcal{Q}$  for any  $g \in G$ , it follows that  $K(\mathcal{Q}) = 0$  where  $K(\cdot)$  is as defined in Lemma 5.1. So that Lemma implies

$$\begin{aligned} F_{\nu \times \mu}^*(\alpha \times_{\sigma} \beta, \mathcal{P} \times \mathcal{Q} | \mathcal{B}_X) &= (1-r)H_{\nu \times \mu}(\mathcal{P} \times \mathcal{Q} | \mathcal{B}_X) + \sum_{i=1}^r h_{\nu \times \mu}((\alpha \times_{\sigma} \beta)_{s_i}, \mathcal{P} \times \mathcal{Q} | \mathcal{B}_X) \\ &= (1-r)H_{\mu}(\mathcal{Q}) + \sum_{i=1}^r h_{\mu}(\beta_{s_i}, \mathcal{Q}) = F_{\mu}^*(\beta, \mathcal{Q}). \end{aligned}$$

□

**Lemma 6.3.** *Let  $\mathcal{Q}$  be a special partition of  $G$ ,  $g \in \Gamma$  and  $\mathcal{P}_g$  denote the partition of  $X$  obtained by pulling  $\beta_g(\mathcal{Q})$  back under the cocycle  $\sigma(g, \cdot)$ . Also, let  $\mathcal{P}'$  be an arbitrary measurable partition of  $X$ . Then*

$$(\alpha \times_{\sigma} \beta)_g((\mathcal{P}_g \vee \mathcal{P}') \times \mathcal{Q}) = \alpha_g(\mathcal{P}_g \vee \mathcal{P}') \times \beta_g(\mathcal{Q})$$

(up to sets of measure zero).

*Proof.* Let  $N$  be the finite-index clopen normal subgroup of  $G$  such that  $\mathcal{Q} = \{qN : q \in G\}$ . Let  $P \in \mathcal{P}_g, P' \in \mathcal{P}'$  and  $qN \in \mathcal{Q}$ . It suffices to show that there exists some  $q'' \in G$  such that

$$(\alpha \times_{\sigma} \beta)_g((P \cap P') \times qN) = \alpha_g(P \cap P') \times q''\beta_g(N)$$

up to sets of measure zero. By definition of  $\mathcal{P}_g$ , there exists a coset  $q'\beta_g(N) \in G/\beta_g(N)$  such that for every  $y \in P$ ,  $\sigma(g, y) \in q'\beta_g(N)$ .

Let  $x \in P \cap P'$  and  $n \in N$ . Then there exists some  $m \in N$  such that

$$(\alpha \times_{\sigma} \beta)_g(x, qn) = (\alpha_g x, \beta_g(qn)\sigma(g, x)) = (\alpha_g x, \beta_g(qn)q'\beta_g(m)).$$

Because  $N$  is normal,  $\beta_g(qn)q'\beta_g(m) \in \beta_g(q)q'\beta_g(N)$ . Thus  $(\alpha \times_{\sigma} \beta)_g(x, qn) \in \alpha_g(P \cap P') \times \beta_g(q)q'\beta_g(N)$ . Since  $(\alpha \times_{\sigma} \beta)_g$  preserves  $\nu \times \mu$ , it follows that  $(\alpha \times_{\sigma} \beta)_g((P \cap P') \times qN) = \alpha_g(P \cap P') \times q''\beta_g(N)$  up to sets of measure zero. □

**Lemma 6.4.** *Let  $\mathcal{P}, \mathcal{Q}$  be measurable partitions for  $\alpha, \beta$  respectively. Suppose  $\mathcal{Q}$  is special and  $\mathcal{P}$  is generating. Let  $\Sigma(\mathcal{P}, \mathcal{Q})$  be the smallest  $\alpha \times_{\sigma} \beta(\Gamma)$ -invariant  $\sigma$ -algebra containing  $\mathcal{P} \times \mathcal{Q}$ . Similarly, let  $\Sigma(\mathcal{Q})$  be the smallest  $\beta(\Gamma)$ -invariant  $\sigma$ -subalgebra of  $\mathcal{B}_G$  which contains  $\mathcal{Q}$ .*

*Then  $\Sigma(\mathcal{P}, \mathcal{Q})$  is the smallest  $\sigma$ -algebra containing  $\mathcal{B}_X \times \Sigma(\mathcal{Q})$  (up to sets of measure zero).*

*Proof.* Clearly,  $\mathcal{P} \times G$  is contained in  $\Sigma(\mathcal{P}, \mathcal{Q})$ . Because

$$(\alpha \times_{\sigma} \beta)_g(\mathcal{P} \times G) = (\alpha_g \mathcal{P}) \times G, \quad \forall g \in \Gamma,$$

it follows that  $(\alpha_g \mathcal{P}) \times G \subset \Sigma(\mathcal{P}, \mathcal{Q})$  for every  $g \in \Gamma$ . Because  $\mathcal{P}$  is generating, this implies  $\mathcal{B}_X \times G \subset \Sigma(\mathcal{P}, \mathcal{Q})$  (up to sets of measure zero).

For each  $g \in \Gamma$ , recall that  $\mathcal{P}_g$  is the partition of  $X$  obtained by pulling  $\beta_g(\mathcal{Q})$  back under the cocycle  $\sigma(g, \cdot)$ . Because  $\sigma(g, \cdot)$  is  $\mathcal{B}_X$ -measurable,  $\mathcal{P}_g \times \mathcal{Q}$  is contained in  $\Sigma(\mathcal{P}, \mathcal{Q})$ . By Lemma 6.3,

$$(\alpha \times_{\sigma} \beta)_g(\mathcal{P}_g \times \mathcal{Q}) = (\alpha_g \mathcal{P}_g) \times (\beta_g \mathcal{Q}) \subset \Sigma(\mathcal{P}, \mathcal{Q})$$

(up to sets of measure zero). Because  $X \times \beta_g \mathcal{Q}$  coarsens  $(\alpha_g \mathcal{P}_g) \times (\beta_g \mathcal{Q})$ , it follows that  $X \times \beta_g \mathcal{Q} \subset \Sigma(\mathcal{P}, \mathcal{Q})$  for every  $g \in \Gamma$ . By definition of  $\Sigma(\mathcal{Q})$ , this implies  $X \times \Sigma(\mathcal{Q}) \subset \Sigma(\mathcal{P}, \mathcal{Q})$ .

Because  $X \times \Sigma(\mathcal{Q})$  and  $\mathcal{B}_X \times G$  generate  $\mathcal{B}_X \times \Sigma(\mathcal{Q})$  (up to sets of measure zero), this implies  $\Sigma(\mathcal{P}, \mathcal{Q}) \supset \mathcal{B}_X \times \Sigma(\mathcal{Q})$ .

To show the opposite inclusion, it suffices to show that  $(\alpha \times_\sigma \beta)_g(\mathcal{P} \times \mathcal{Q}) \in \mathcal{B}_X \times \Sigma(\mathcal{Q})$  for any  $g \in \Gamma$ . By the previous lemma,

$$(\alpha \times_\sigma \beta)_g(\mathcal{P} \times \mathcal{Q}) \leq (\alpha \times_\sigma \beta)_g((\mathcal{P}_g \vee \mathcal{P}) \times \mathcal{Q}) = (\alpha_g(\mathcal{P}_g \vee \mathcal{P})) \times (\beta_g \mathcal{Q}) \in \mathcal{B}_X \times \Sigma(\mathcal{Q}).$$

□

*Proof of Theorem 3.3.* Let  $\mathcal{P}$  be a finite-entropy generating partition for  $\alpha$  and  $\mathcal{Q}$  be a special generating partition for  $\beta$ . By the previous lemma,  $\mathcal{P} \times \mathcal{Q}$  is generating for  $\alpha \times_\sigma \beta$ . So Theorems 4.1 and 4.2 imply

$$\begin{aligned} f_{\nu \times \mu}(\alpha \times_\sigma \beta) - f_\nu(\alpha) &= f_{\nu \times \mu}(\alpha \times_\sigma \beta | \mathcal{B}_X) \\ &= \inf_{n > 0} F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, (\mathcal{P} \times \mathcal{Q})^{B(n)} | \mathcal{B}_X). \end{aligned}$$

For each  $g \in \Gamma$ , let  $\mathcal{P}_g$  be the partition of  $X$  obtained by pulling  $(\beta_g \mathcal{Q})$  back under  $\sigma(g, \cdot)$ . By Lemma 6.3, for any partition  $\mathcal{P}'$  of  $X$

$$(\alpha \times_\sigma \beta)_g((\mathcal{P}_g \vee \mathcal{P}') \times \mathcal{Q}) = (\alpha_g(\mathcal{P}_g \vee \mathcal{P}')) \times (\beta_g \mathcal{Q}).$$

Given an integer  $n > 0$  let  $\mathcal{R}_n = \bigvee_{g \in B(n)} \mathcal{P}_g$ . By Lemma 6.3,

$$\begin{aligned} (\mathcal{P} \vee \mathcal{R}_n) \times \mathcal{Q}^{B(n)} &= \bigvee_{w \in B(n)} (\alpha \times_\sigma \beta)_w((\mathcal{P} \vee \mathcal{R}_n) \times \mathcal{Q}) \\ &= \bigvee_{w \in B(n)} \alpha_w(\mathcal{P} \vee \mathcal{R}_n) \times \beta_w(\mathcal{Q}) \\ &= (\mathcal{P} \vee \mathcal{R}_n)^{B(n)} \times \mathcal{Q}^{B(n)}. \end{aligned}$$

Because we are conditioning on  $\mathcal{B}_X$  and  $(\mathcal{R}_n \times G)^{B(n)} = (\mathcal{R}_n^{B(n)} \times G)$ ,

$$\begin{aligned} F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, (\mathcal{P} \times \mathcal{Q})^{B(n)} | \mathcal{B}_X) &= F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, (\mathcal{P} \times \mathcal{Q})^{B(n)} \vee \mathcal{R}_n^{B(n)} \times G | \mathcal{B}_X) \\ &= F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, ((\mathcal{P} \vee \mathcal{R}_n) \times \mathcal{Q})^{B(n)} | \mathcal{B}_X) \\ &= F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, ((\mathcal{P} \vee \mathcal{R}_n)^{B(n)} \times \mathcal{Q}^{B(n)}) | \mathcal{B}_X). \end{aligned}$$

By Lemma 6.2,

$$F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, ((\mathcal{P} \vee \mathcal{R}_n)^{B(n)} \times \mathcal{Q}^{B(n)}) | \mathcal{B}_X) = F_\mu^*(\beta, \mathcal{Q}^{B(n)}).$$

So we now have

$$\begin{aligned} f_{\nu \times \mu}(\alpha \times_\sigma \beta) &= f_\nu(\alpha) + f_{\nu \times \mu}(\alpha \times_\sigma \beta | \mathcal{B}_X) \\ &= f_\nu(\alpha) + \inf_{n > 0} F_{\nu \times \mu}^*(\alpha \times_\sigma \beta, (\mathcal{P} \times \mathcal{Q})^{B(n)} | \mathcal{B}_X) \\ &= f_\nu(\alpha) + \inf_{n > 0} F_\mu^*(\beta, \mathcal{Q}^{B(n)}) \\ &= f_\nu(\alpha) + f_\mu(\beta). \end{aligned}$$

The last equality holds by Theorem 2.1. □

## 7. EXAMPLES

It is convenient to introduce the following notation. Let  $\Gamma = \langle s_1, \dots, s_r \rangle$  be the rank  $r$  free group. If  $K$  is a set then  $K^\Gamma$  is the set of all functions  $x : \Gamma \rightarrow K$ . The *shift-action* of  $\Gamma$  on  $K^\Gamma$  is defined as follows. For  $g, f \in \Gamma$  and  $x \in K^\Gamma$ ,  $gx \in K^\Gamma$  is the map  $(gx)(f) = x(g^{-1}f)$ .

If  $\Gamma$  acts on a compact group  $G$  and the action is understood, we write  $f(\Gamma \curvearrowright G)$  to mean the  $f$ -invariant of the action of  $G$  with respect to Haar measure.

**7.1. The Ornstein-Weiss Example.** This example comes from the appendix to [OW87]. To explain its relevance, let us recall some basic facts from classical entropy theory. Let  $\Delta$  be a countable amenable group,  $K$  a finite set and  $u$  the uniform probability measure on  $K$ . It is straightforward to compute the entropy of the shift action of  $\Delta$  on  $(K^\Delta, u^\Delta)$ : it is  $\log |K|$ . Because entropy never increases under a factor map, it follows that if  $|K| > 1$  then the action  $\Delta \curvearrowright (K^\Delta, u^\Delta)$  cannot factor onto the action  $\Delta \curvearrowright ((K \times K)^\Delta, (u \times u)^\Delta)$ .

By contrast, Ornstein and Weiss showed that if  $\Gamma$  is the rank 2 free group then  $\Gamma \curvearrowright (\mathbb{Z}/2\mathbb{Z})^\Gamma$  factors onto  $\Gamma \curvearrowright (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})^\Gamma$ . This convinced many researchers that there could not be an entropy theory for free groups.

The factor map is defined by

$$\phi : (\mathbb{Z}/2\mathbb{Z})^\Gamma \rightarrow (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})^\Gamma,$$

$$\phi(x)(g) = (x(g) + x(gs_1), x(g) + x(gs_2)), \forall x \in (\mathbb{Z}/2\mathbb{Z})^\Gamma, g \in \Gamma.$$

We consider  $(\mathbb{Z}/2\mathbb{Z})^\Gamma$  and  $(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})^\Gamma$  as compact groups under pointwise addition. It is a straightforward exercise to show that  $\phi$  is a surjective homomorphism which is equivariant with respect to the shift-actions of  $\Gamma$  and therefore, defines a factor map. Moreover, the kernel of  $\phi$  consists of two elements,  $x_0, x_1$ , where  $x_i : \Gamma \rightarrow \mathbb{Z}/2\mathbb{Z}$  is defined by  $x_i(g) = i$ . Let  $N = \{x_0, x_1\}$ . Because  $N$  is finite, it clearly satisfies the conditions of Theorem 2.2. So that result implies

$$f(\Gamma \curvearrowright (\mathbb{Z}/2\mathbb{Z})^\Gamma) = f(\Gamma \curvearrowright N) + f(\Gamma \curvearrowright (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})^\Gamma).$$

In [Bo10a], it is shown that  $f(\Gamma \curvearrowright (\mathbb{Z}/2\mathbb{Z})^\Gamma) = \log(2)$  and  $f(\Gamma \curvearrowright (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})^\Gamma) = \log(4)$  as expected. Therefore,  $f(\Gamma \curvearrowright N) = -\log(2)$ . This is easy to verify by direct computation.

**7.2. A generalization.** The example above can be generalized with the help of [MRV11, proof of Theorem B] which states the following: if  $\Gamma = \langle s_1, \dots, s_r \rangle$  is any finite rank free group,  $K$  is any compact second countable group,  $K^\Gamma$  is the group of all functions  $x : \Gamma \rightarrow K$  under pointwise multiplication and  $K$  is identified with the constant functions in  $K^\Gamma$  then the action  $\Gamma \curvearrowright K^\Gamma/K$  is measurably conjugate to  $\Gamma \curvearrowright (K^r)^\Gamma$  (where the measures involved are the Haar measures and the actions are the shift actions).

When  $K$  is finite, we can apply Theorem 2.2 to obtain

$$f(\Gamma \curvearrowright K^\Gamma) = f(\Gamma \curvearrowright K) + f(\Gamma \curvearrowright (K^r)^\Gamma).$$

This is easy to check:  $f(\Gamma \curvearrowright K^\Gamma) = \log(|K|)$  and  $f(\Gamma \curvearrowright (K^r)^\Gamma) = r \log(|K|)$  by [Bo10a]. By a straightforward computation,  $f(\Gamma \curvearrowright K) = -(r-1) \log |K|$ .

**7.3. An algebraic example.** As above, let  $\Gamma = \langle s_1, \dots, s_r \rangle$  be a finite rank free group. Let  $p > 1$  be a prime number and  $h \in (\mathbb{Z}/p\mathbb{Z})^\Gamma$ . We consider  $h$  as a function from  $\Gamma$  to  $\mathbb{Z}$  such that  $h(s) = 0$  for all but finitely many  $s \in \Gamma$ . Define the convolution operator  $\phi_h : (\mathbb{Z}/p\mathbb{Z})^\Gamma \rightarrow (\mathbb{Z}/p\mathbb{Z})^\Gamma$  by

$$\phi_h(x)(g) = \sum_{s \in \Gamma} x(gs)h(s^{-1}), \quad \forall g \in \Gamma.$$

This is a  $\Gamma$ -equivariant homomorphism. Let  $X_{h,p}$  denote the kernel of  $\phi_h$ . Let  $X_{h,p}^* < X_{h,p}$  be the subgroup consisting of all elements  $x \in X_{h,p}$  with  $x(e) = 0$ . This is a finite-index normal clopen subgroup and  $\{gX_{h,p}^* : g \in X_{h,p}\}$  is a generating partition for the shift-action of  $\Gamma$ . Therefore, we can apply Theorem 2.2 to obtain

$$f(\Gamma \curvearrowright (\mathbb{Z}/p\mathbb{Z})^\Gamma) = f(\Gamma \curvearrowright X_{h,p}) + f(\Gamma \curvearrowright \phi_h((\mathbb{Z}/p\mathbb{Z})^\Gamma)).$$

**Theorem 7.1.**  $\phi_h$  is onto if  $h$  is nonzero.

Therefore,

$$f(\Gamma \curvearrowright \phi_h((\mathbb{Z}/p\mathbb{Z})^\Gamma)) = f(\Gamma \curvearrowright (\mathbb{Z}/p\mathbb{Z})^\Gamma).$$

Thus  $f(\Gamma \curvearrowright X_{h,p}) = 0$ .

To prove Theorem 7.1, we need a little preparation.

**Definition 7.2.** Let  $C_r$  be the Cayley graph of  $\Gamma$ . It has vertex set  $\Gamma$  and edges  $\{g, gs_i\}$  for all  $g \in \Gamma$  and  $1 \leq i \leq r$ . Given a set  $F \subset \Gamma$ , the *induced subgraph* of  $F$  is the subgraph  $C_r(F) \subset C_r$  which has vertex set  $F$  and contains every edge of  $C_r$  which has both endpoints in  $F$ . A subset  $F \subset \Gamma$  is said to be *connected* if its induced subgraph in  $C_r$  is connected. The *convex hull* of a set  $F \subset \Gamma$  is the smallest connected set  $F' \subset \Gamma$  with  $F \subset F'$ . An *extreme point* of  $F$  is an element  $f \in F$  that has degree 1 in  $C_r(F)$ . We let  $\text{Ex}(F)$  denote the set of extreme points of  $F$ . Note that if  $F'$  is the convex hull of  $F$  then  $\text{Ex}(F') \subset F$ .

**Lemma 7.3.** Let  $F = \{g \in \Gamma : h(g^{-1}) \neq p\mathbb{Z}\}$ . Let  $\overline{F}$  be the convex hull of  $F$ . Suppose there exists an ordering  $\gamma_0, \gamma_1, \gamma_2, \dots$  of  $\Gamma$  such that for every  $n \geq 1$   $\{\gamma_0, \dots, \gamma_n\}$  is connected and

$$\gamma_n \overline{F} \not\subseteq \cup_{i=0}^{n-1} \gamma_i \overline{F}.$$

Then  $\phi_h$  is onto.

*Proof.* By compactness of  $(\mathbb{Z}/p\mathbb{Z})^\Gamma$  and continuity of  $\phi_h$ , it suffices to show that for every  $y \in (\mathbb{Z}/p\mathbb{Z})^\Gamma$  and every  $n \geq 0$ , there exists an  $x \in (\mathbb{Z}/p\mathbb{Z})^\Gamma$  such that  $\phi_h(x)(\gamma_i) = y(\gamma_i)$  for every  $0 \leq i \leq n$ . We will prove this statement by induction on  $n$ . It is clearly true for  $n = 0$ . So suppose there is an  $n \geq 0$  for which the statement is true. Fix  $y \in (\mathbb{Z}/p\mathbb{Z})^\Gamma$  and let  $x \in (\mathbb{Z}/p\mathbb{Z})^\Gamma$  be such that  $\phi_h(x)(\gamma_i) = y(\gamma_i)$  for every  $0 \leq i \leq n$ .

By hypothesis,  $\gamma_{n+1} \overline{F} \not\subseteq \cup_{i=0}^n \gamma_i \overline{F}$ . Because  $\cup_{i=0}^n \gamma_i \overline{F}$  and  $\gamma_{n+1} \overline{F}$  are connected and the convex hull of the extreme points set of a connected set is the connected set itself, there must be an extremal point  $f \in \text{Ex}(\overline{F})$  such that  $\gamma_{n+1} f \notin \cup_{i=0}^n \gamma_i \overline{F}$ . However,  $\text{Ex}(\overline{F}) \subset F$ . So  $f \in F$ . By definition, this means that  $h(f^{-1}) \neq p\mathbb{Z}$ . Because  $p$  is prime, we may therefore define an element  $m \in \mathbb{Z}/p\mathbb{Z}$  by

$$m = h(f^{-1})^{-1} \left( y(\gamma_{n+1}) - \sum_{g \in \Gamma \setminus \{f\}} x(\gamma_{n+1}g)h(g^{-1}) \right).$$

Define  $x' \in (\mathbb{Z}/p\mathbb{Z})^\Gamma$  by  $x'(g) = x(g)$  if  $g \neq \gamma_{n+1}f$  and  $x'(\gamma_{n+1}f) = m$ . Because  $\gamma_{n+1}f \notin \cup_{i=0}^n \gamma_i \overline{F}$ , it follows that  $\phi_h(x')(\gamma_i) = \phi_h(x)(\gamma_i)$  for all  $0 \leq i \leq n$ . Also a straightforward computation shows  $\phi_h(x')(\gamma_{n+1}) = y(\gamma_{n+1})$ . So  $\phi_h(x')(\gamma_i) = y(\gamma_i)$  for all  $0 \leq i \leq n+1$ . This completes the inductive step and the claim.  $\square$

**Definition 7.4.** Let  $S = \{s_1, \dots, s_r\}$ . For  $g \in \Gamma$ , let  $|g|$  be the smallest number  $n \geq 0$  such that there exist elements  $t_1, \dots, t_n \in S \cup S^{-1}$  with  $g = t_1 \cdots t_n$ . We also let  $d(g_1, g_2) = |g_1^{-1}g_2|$  for any  $g_1, g_2 \in \Gamma$ . For  $g \in \Gamma$  and  $n \geq 0$ , let  $B(g, n) = \{k \in \Gamma : d(k, g) \leq n\}$  be the ball of radius  $n$  centered at  $g$ .

Let  $K \subset \Gamma$  be a finite set. The *radius* of  $K$  is the smallest number  $r \geq 0$  such that there exists a  $v \in \Gamma$  such that  $B(v, r) \supset K$ . An element  $v \in \Gamma$  is called a *center* of  $K$  if  $B(v, r) \supset K$  where  $r$  is the radius of  $K$ . For any  $v, w \in \Gamma$ , we let  $[v, w] \subset \Gamma$  be the set of all  $g \in \Gamma$  such that the shortest path from  $v$  to  $w$  in the Cayley graph  $C_r$  contains  $g$ .

**Lemma 7.5.** *Let  $K$  be a connected finite set with radius  $r \geq 1$ . Suppose the identity element  $e$  is a center of  $K$ . Then there exist elements  $v, w \in K$  such that  $[e, v] \cap [e, w] = \{e\}$ ,  $|v| = r$  and  $|w| \in \{r-1, r\}$ .*

*Proof.* Because  $K$  has radius  $r$  and center  $e$ , there is an element  $v$  with  $|v| = r$ . To obtain a contradiction, suppose that there is no  $w \in K$  with  $|w| \in \{r-1, r\}$  and  $[e, v] \cap [e, w] = \{e\}$ . Let  $v_1 \in S \cup S^{-1}$  be the unique element with  $|v_1^{-1}v| = r-1$ . We claim that  $B(v_1, r-1) \supset K$ . To see this, let  $w \in K$ . If  $|w| \leq r-2$  then  $w \in B(e, r-2) \subset B(v_1, r-1)$ . If  $|w| > r-2$  then, because  $K$  has center  $e$  and radius  $r$ ,  $|w| \in \{r-1, r\}$ . By assumption, this implies  $[e, v] \cap [e, w] \neq \{e\}$ . So let  $y \in [e, v] \cap [e, w]$  with  $y \neq e$ . Then  $[e, y] \subset [e, v]$ . This implies that  $v_1 \in [e, y]$ . In particular,  $v_1 \in [e, v] \cap [e, w]$ , so  $v_1 \in [e, w]$ . Because  $|w| \leq r$ , this implies  $d(v_1, w) \leq r-1$  as claimed. So we have shown that in all cases, if  $w \in K$  then  $w \in B(v_1, r-1)$ . This shows that the radius of  $K$  is at most  $r-1$ , a contradiction. This contradiction proves the lemma.  $\square$

**Lemma 7.6.** *Let  $K$  be a connected finite set with radius  $r \geq 1$ . Suppose the identity element  $e$  is a center of  $K$ . Suppose  $g_1, \dots, g_n \in \Gamma \setminus \{e\}$  are elements with*

$$K \subset \cup_{i=1}^n g_n K.$$

*Then  $e$  is contained in the convex hull of  $\{g_1, \dots, g_n\}$ .*

*Proof.* Let  $v, w \in K$  be elements such that  $[e, v] \cap [e, w] = \{e\}$ ,  $|v| = r$  and  $|w| \in \{r-1, r\}$ . Let  $g_i, g_j \in \{g_1, \dots, g_n\}$  be such that  $v \in g_i K$  and  $w \in g_j K$ . Let  $x, y \in K$  be such that  $v = g_i x$  and  $w = g_j y$ .

Let  $v_1, v_2, x_1, x_2 \in \Gamma$  be such that  $v = v_1 v_2$ ,  $|v| = |v_1| + |v_2|$ ,  $x_2 = v_2$ ,  $x = x_1 x_2$ ,  $|x| = |x_1| + |x_2|$  and  $|v_2| = |x_2|$  is as large as possible. Thus  $g_i = v x^{-1} = v_1 x_1^{-1}$  and  $|v x^{-1}| = |v_1| + |x_1|$ . Because  $r$  is the radius of  $K$ ,  $e$  is a center and  $x \in K$  we have  $|x| \leq r$ . Also, we cannot have  $v = x$  (since this would imply  $g_i = v x^{-1} = e$ , a contradiction). So we must have  $|v_1| \geq 1$ . Thus  $[e, v] \cap [e, g_i] \neq \{e\}$ .

Let  $w_1, w_2, y_1, y_2 \in \Gamma$  be such that  $w = w_1 w_2$ ,  $|w| = |w_1| + |w_2|$ ,  $y_2 = w_2$ ,  $y = y_1 y_2$ ,  $|y| = |y_1| + |y_2|$  and  $|w_2| = |y_2|$  is as large as possible. Thus  $g_j = w y^{-1} = w_1 y_1^{-1}$  and  $|w y^{-1}| = |w_1| + |y_1|$ . Because  $r$  is the radius of  $K$ ,  $e$  is a center and  $y \in K$  we have  $|y| \leq r$ .

**Case 1.** If  $|w| = r$ , then, as in the previous paragraph, we must have  $[e, w] \cap [e, g_j] \neq \{e\}$ . Because  $[e, v] \cap [e, w] = \{e\}$ , this implies  $e \in [g_i, g_j]$  which implies the lemma.

**Case 2.** Suppose  $|w| = r - 1$  and  $|w_1| \geq 1$ . Thus  $[e, w] \cap [e, g_j] \neq \{e\}$ . Because  $[e, v] \cap [e, w] = \{e\}$ , this implies  $e \in [g_i, g_j]$  which implies the lemma.

**Case 3.** Suppose  $|w| = r - 1$  and  $|w_1| = 0$ . Then  $w = w_2$ , so  $|w_2| = r - 1$ . Because  $g_j = wy^{-1} = w_1y_1^{-1} = y_1^{-1} \neq e$ , we must  $y_1 \neq e$ . Thus  $|y| = |y_1| + |y_2| = |y_1| + |w_2| = |y_1| + r - 1$ . Because  $y \in K$  and  $K$  has radius  $r$  and center  $e$ , we must have  $|y_1| = 1$  and  $|y| = r$ . If  $[e, y] \cap [e, v] = \{e\}$  then, after replacing  $w$  with  $y$  we are in the situation of Case 1 (note  $y = g_k y'$  for some  $1 \leq k \leq n$  and  $y' \in K$ ). So we may assume  $[e, y] \cap [e, v] \neq \{e\}$  which implies  $y_1 \in [e, v]$ . Because  $g_j = y_1^{-1}$ , and  $[e, v] \cap [e, g_i] \neq \{e\}$ , we have  $[e, g_i] \cap [e, g_j] = \{e\}$  which implies  $e \in [g_i, g_j]$  which implies the lemma.

[Proof of Theorem 7.1] Let  $F = \{g \in \Gamma : h(g^{-1}) \neq p\mathbb{Z}\}$ . Let  $\overline{F}$  be the convex hull of  $F$ . For any  $g \in \Gamma$ ,  $\phi_h$  is onto if and only if  $\phi_{gh}$  is onto. So after replacing  $h$  with  $gh$  for some  $g \in \Gamma$ , we may assume that  $e$  is a center of  $\overline{F}$ .

Let  $g_0, g_1, \dots$  be an ordering of  $\Gamma$  such that for every  $n \geq 0$ ,  $\{g_0, \dots, g_n\}$  is connected. We claim that for every  $n \geq 1$ ,

$$\gamma_n \overline{F} \not\subseteq \cup_{i=0}^{n-1} \gamma_i \overline{F}.$$

To obtain a contradiction, suppose that the claim is false for some  $n \geq 1$ . Then  $\overline{F} \subset \cup_{i=0}^{n-1} \gamma_n^{-1} \gamma_i \overline{F}$ ,  $\gamma_n^{-1} \gamma_i \neq e$  for any  $0 \leq i \leq n-1$  and because  $\{\gamma_0, \dots, \gamma_{n-1}\}$  is connected,  $\{\gamma_n^{-1} \gamma_0, \dots, \gamma_n^{-1} \gamma_{n-1}\}$  is connected which implies that  $e$  is not in the convex hull of  $\{\gamma_n^{-1} \gamma_0, \dots, \gamma_n^{-1} \gamma_{n-1}\}$ . This contradicts the previous lemma. So we must have that for every  $n \geq 1$ ,

$$\gamma_n \overline{F} \not\subseteq \cup_{i=0}^{n-1} \gamma_i \overline{F}.$$

The theorem now follows from Lemma 7.3. □

## REFERENCES

- [BG12] L. Bowen and Y. Gutman. *Corrigendum to “Nonabelian free group actions: Markov processes, the Abramov-Rohlin formula and Yuzvinskii’s formula”*. To appear in *Ergodic Theory Dyn. Syst.* <http://arxiv.org/abs/1206.6772>. 2012.
- [BM09] M. Björklund and R. Miles. *Entropy range problems and actions of locally normal groups*. *Discrete Contin. Dyn. Syst.* 25 (2009), no. 3, 981–989.
- [Bo10a] L. Bowen. *A measure-conjugacy invariant for actions of free groups*. *Ann. of Math. (2)* **171** (2010), no. 2, 1387–1400.
- [Bo10b] L. Bowen. *Measure conjugacy invariants for actions of countable sofic groups*. *J. Amer. Math. Soc.* 23 (2010), 217–245.
- [Bo10c] L. Bowen. *Nonabelian free group actions: Markov processes, the Abramov-Rohlin formula and Yuzvinskii’s formula*. *Ergodic Theory Dynam. Systems* 30 (2010), no. 6, 1629–1663.
- [He01] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*. Corrected reprint of the 1978 original. *Graduate Studies in Mathematics*, 34. American Mathematical Society, Providence, RI, 2001. xxvi+641 pp.
- [Ju65] S. A. Juzvinskii. *Metric properties of the endomorphisms of compact groups*. *Izv. Akad. Nauk SSSR Ser. Mat.* 29 1965 1295–1328.
- [KH95] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems, with a supplement by Anatole Katok and Leonardo Mendoza*. *Encyclopedia of Mathematics and its Applications* 54, Cambridge University Press, 1995, 822 pp.
- [Ku65] A. G. Kushnirenko. *An upper bound of the entropy of classical dynamical systems*. *Sov. Math. Dokl.* 6 (1965), 360–362.
- [Li11] H. Li. *Compact group automorphisms, addition formulas and Fuglede-Kadison determinants*. arXiv:1001.0419.

- [LS09] D. Lind and K. Schmidt. Preprint. 2009.
- [LSW90] D. Lind, K. Schmidt and T. Ward. *Mahler measure and entropy for commuting automorphisms of compact groups*. Invent. Math. 101 (1990), no. 3, 593–629.
- [MRV11] N. Meesschaert, S. Raum and S. Vaes. *Stable orbit equivalence of Bernoulli actions of free groups and isomorphism of some of their factor actions*. arXiv:1107.1357
- [Mi08] R. Miles. *The entropy of algebraic actions of countable torsion-free abelian groups*. Fund. Math. 201 (2008), 261D282.
- [OW87] D. Ornstein and B. Weiss. *Entropy and isomorphism theorems for actions of amenable groups*. J. Analyse Math. 48 (1987), 1–141.
- [Th71] R. K. Thomas. *The addition theorem for the entropy of transformations of  $G$ -spaces*. Trans. Amer. Math. Soc. 160 (1971), 119–130.

Lewis Bowen, Mathematics Department, Mailstop 3368, Texas A&M University College Station, TX 77843-3368 United States.

*E-mail address:* `lpbowen@math.tamu.edu`

Yonatan Gutman, Institute of Mathematics, Polish Academy of Sciences, ul. Śniadeckich 8, 00-956 Warszawa, Poland.

*E-mail address:* `y.gutman@impan.pl`